arXiv:1208.0589v1 [math.NT] 2 Aug 2012

DIRICHLET'S TYPE OF TWISTED EULERIAN POLYNOMIALS IN CONNECTION WITH TWISTED EULERIAN-$\mathcal{L}$-FUNCTION

SERKAN ARACI AND **MEHMET ACIKGOZ**

Abstract. In the present paper, we effect Dirichlet's type of twisted Eulerian polynomials by using p-adic fermionic q-integral on the p-adic integer ring. Also, we introduce some new interesting identities for them. As a result of them, by using contour integral on the generating function of Dirichlet's type of twisted Eulerian polynomials and so we define twisted Eulerian-$\mathcal{L}$-function which interpolates of Dirichlet's type of Eulerian polynomials at negative integers which we state in this paper.

2010 Mathematics Subject Classification. Primary 05A10, 11B65; Secondary 11B68, 11B73.

Keywords and phrases. Eulerian polynomials, p-adic fermionic q-integral on $\mathbb{Z}_p$, contour integral, L-function, Dirichlet's character.

1. Introduction

As it is well-known, Eulerian polynomials, $\mathcal{A}_n(x)$, are given by means of the following exponential generating function:

$$e^{\mathcal{A}(x)t}=\sum_{n=0}^{\infty}\mathcal{A}_n(x)\frac{t^n}{n!}=\frac{1-x}{e^{t(1-x)}-x} \tag{1}$$

where $\mathcal{A}^n(x):=\mathcal{A}_n(x)$, symbolically. Eulerian polynomials can find via the following recurrence relation:

$$(\mathcal{A}(t)+(t-1))^n-t\mathcal{A}_n(t)=\begin{cases}1-t & \text{if } n=0\\ 0 & \text{if } n\neq 0,\end{cases} \tag{2}$$

(for details, see [2], [13] and [22]).

Imagine that p be a fixed odd prime number. Throughout this paper, we use the following notations. By $\mathbb{Z}_p$, we denote the ring of p-adic rational integers, $\mathbb{Q}$ denotes the field of rational numbers, $\mathbb{Q}_p$ denotes the field of p-adic rational numbers, and $\mathbb{C}_p$ denotes the completion of algebraic closure of $\mathbb{Q}_p$. Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{N}^*=\mathbb{N}\cup\{0\}$.

The p-adic absolute value is defined by

$$|p|_p=\frac{1}{p}.$$

Let q be an indeterminate with $|q-1|_p<1$. Thus, we give definition of q-integer of x, which is defined by

$$[x]_q=\frac{1-q^x}{1-q} \text{ and } [x]_{-q}=\frac{1-(-q)^x}{1+q},$$

where we note that $\lim_{q\to1}[x]_q=x$ (see [1-26]).

Let $UD\left(\mathbb{Z}_p\right)$ be the space of uniformly differentiable functions on $\mathbb{Z}_p$. For a positive integer d with $(d,p)=1$, let

$$X=X_d=\lim_{\overleftarrow{n}}\mathbb{Z}/dp^n\mathbb{Z}=\overset{dp-1}{\underset{a=0}{\cup}}\left(a+dp\mathbb{Z}_p\right)$$

with

$$a+dp^n\mathbb{Z}_p=\left\{x\in X\mid x\equiv a\left(\operatorname{mod}dp^n\right)\right\}$$

where $a\in\mathbb{Z}$ satisfies the condition $0\leq a<dp^n$ and let $\sigma:X\rightarrow\mathbb{Z}_p$ be the transformation introduced by the inverse limit of the natural transformation

$$\mathbb{Z}/dp^n\mathbb{Z}\mapsto\mathbb{Z}/p^n\mathbb{Z}.$$

If f is a function on $\mathbb{Z}_p$, then we will utilize the same notation to indicate the function $f\circ\sigma$.

For a continuous function $f:X\rightarrow\mathbb{C}_p$, the p-adic q-integral on $\mathbb{Z}_p$ is defined by T. Kim in [3] and [4], as follows:

$$I_q\left(f\right)=\int_X f\left(\upsilon\right)d\mu_q\left(\upsilon\right)=\int_{\mathbb{Z}_p}f\left(\upsilon\right)d\mu_q\left(\upsilon\right)=\lim_{n\rightarrow\infty}\frac{1}{\left[p^n\right]_q}\sum_{\upsilon=0}^{p^n-1}q^{\upsilon}f\left(\upsilon\right).\tag{3}$$

The bosonic integral is considered as the bosonic limit $q\rightarrow1$, $I_1\left(f\right)=\lim_{q\rightarrow1}I_q\left(f\right)$. In [8], [9] and [10], similarly, the p-adic fermionic integration on $\mathbb{Z}_p$ is given by Kim as follows:

$$I_{-q}\left(f\right)=\lim_{q\rightarrow-q}I_q\left(f\right)=\int_{\mathbb{Z}_p}f\left(\upsilon\right)d\mu_{-q}\left(\upsilon\right).\tag{4}$$

By (4), we have the following well-known integral equation:

$$q^nI_{-q}\left(f_n\right)+\left(-1\right)^{n-1}I_{-q}\left(f\right)=\left[2\right]_q\sum_{l=0}^{n-1}\left(-1\right)^{n-1-l}q^lf\left(l\right)\tag{5}$$

here let $f_n\left(x\right)$ be a translation with $f_n\left(x\right):=f\left(x+n\right)$. By (5), we readily derive the following

$$qI_{-q}\left(f_1\right)+I_{-q}\left(f\right)=\left[2\right]_qf\left(0\right).\tag{6}$$

Replacing q by q^{-1} in (6), we easily derive the following

$$I_{-q^{-1}}\left(f_1\right)+qI_{-q^{-1}}\left(f\right)=\left[2\right]_qf\left(0\right).\tag{7}$$

Recently, Kim *et al.* [2] is considered $f(x)=e^{-x(1+q)t}$ in (7), then they gave Witt's formula of Eulerian polynomials as follows:

For $n\in\mathbb{N}^*$,

$$I_{-q^{-1}}\left(x^n\right)=\frac{\left(-1\right)^n}{\left(1+q\right)^n}\mathcal{A}_n\left(-q\right).\tag{8}$$

In previous paper [22], Araci, Acikgoz and Gao are introduced generating function of Dirichlet's type of Eulerian polynomials as

$$\sum_{n=0}^{\infty}\mathcal{A}_{n,\chi}\left(-q\right)\frac{t^n}{n!}=\left[2\right]_q\sum_{l=0}^{d-1}\left(-1\right)^lq^{d-l+1}\chi\left(l\right)\frac{e^{-l(1+q)t}}{e^{-d(1+q)t}+q^d}.\tag{9}$$

Also, they gave Witt's formula for Dirichlet's type of Eulerian polynomials, as follows:

$$I_{-q^{-1}}\left(\chi\left(x\right)x^{n}\right)=\frac{\left(-1\right)^{n}}{\left(1+q\right)^{n}}\mathcal{A}_{n,\chi}\left(-q\right).$$

In this paper, we will also consider Dirichlet's type of twisted Eulerian polynomials. By applying Mellin transformation to generating function of Dirichlet's type of twisted Eulerian polynomials, we will describe twisted Eulerian-$\mathcal{L}$-function. Next, by utilizing from Cauchy-Residue theorem, we will see that twisted Eulerian-$\mathcal{L}$-function is related to Dirichlet's type of twisted Eulerian polynomials at negative integers.

2. On the Dirichlet's type of twisted Eulerian polynomials

By using (5), it is easy to see that

$$I_{-q^{-1}}\left(f_{d}\right)+q^{d}I_{-q^{-1}}\left(f\right)=\left[2\right]_{q}\sum_{l=0}^{d-1}\left(-1\right)^{l}q^{d-l+1}f\left(l\right). \tag{10}$$

Let $C_{p^n}=\left\{\zeta\mid\zeta^{p^n}=1\right\}$ be the cyclic group of order p^n, and let

$$\mathcal{T}_p=\lim_{n\to\infty}C_{p^n}=C_{p^\infty}=\underset{n\geq 0}{\cup}C_{p^n},$$

we want to note that $\mathcal{T}_p$ is locally constant space.

Let χ be a Dirichlet's character of conductor $d\,(=\text{odd})$ and $\zeta\in\mathcal{T}_p$, then, taking $f(x)=\zeta^{x}\chi\left(x\right)e^{-x(1+q)t}$ in (10), then it is equality to

$$\begin{aligned}I_{-q^{-1}}&\left(\zeta^{x+d}\chi\left(x+d\right)e^{-(x+d)(1+q)t}\right)+q^{d}I_{-q^{-1}}\left(\zeta^{x}\chi\left(x\right)e^{-x(1+q)t}\right)\\&=\left[2\right]_{q}\sum_{l=0}^{d-1}\left(-1\right)^{l}q^{d-l+1}\chi\left(l\right)e^{-l(1+q)t}\zeta^{l}\end{aligned}$$

From this, we easily see that

$$I_{-q^{-1}}\left(\zeta^{x}\chi\left(x\right)e^{-x(1+q)t}\right)=\left[2\right]_{q}\sum_{l=0}^{d-1}\left(-1\right)^{l}q^{d-l+1}\zeta^{l}\chi\left(l\right)\frac{e^{-l(1+q)t}}{\zeta^{d}e^{-d(1+q)t}+q^{d}}. \tag{11}$$

Now, we give definition of generating function of Dirichlet's type of twisted Eulerian polynomials as follows:

Definition 1. *Let* $\mathcal{G}_{q,\zeta}\left(t\mid\chi\right)=\sum_{n=0}^{\infty}\mathcal{A}_{n,\chi,\zeta}\left(-q\right)\frac{t^n}{n!}$ *and* $\zeta\in T_p$. *Then, we define twisted Dirichlet's type of Eulerian polynomials by means of the following generating function:*

$$\mathcal{G}_{q,\zeta}\left(t\mid\chi\right)=\left[2\right]_{q}\sum_{l=0}^{d-1}\left(-1\right)^{l}q^{d-l+1}\zeta^{l}\chi\left(l\right)\frac{e^{-l(1+q)t}}{\zeta^{d}e^{-d(1+q)t}+q^{d}}.$$

By considering the above definition, we readily derive the following corollary.

Corollary 2.1. *For any* $\zeta\in\mathcal{T}_p$ *and* $n\in\mathbb{N}^*$, *then we have*

$$\mathcal{A}_{n,\chi,\zeta}\left(-q\right)=\textit{coefficient of }\frac{t^n}{n!}\textit{ of }\left[2\right]_{q}\sum_{l=0}^{d-1}\left(-1\right)^{l}q^{d-l+1}\zeta^{l}\chi\left(l\right)\frac{e^{-l(1+q)t}}{\zeta^{d}e^{-d(1+q)t}+q^{d}}.$$

By using (11) and Definition 1, we procure the following theorem which is the Witt's formula for Dirichlet's type of twisted Eulerian polynomials.

Theorem 2.2. *The following equality*

$$I_{-q^{-1}}\left(\zeta^{x}\chi\left(x\right)x^{n}\right)=\int_{\mathbb{Z}_p}\zeta^{x}\chi\left(x\right)x^{n}d\mu_{-q^{-1}}\left(x\right)=\frac{\left(-1\right)^{n}}{\left(1+q\right)^{n}}\mathcal{A}_{n,\chi,\zeta}\left(-q\right) \tag{12}$$

holds true.

By using Definition 1, becomes

$$\begin{aligned}\sum_{n=0}^{\infty}\mathcal{A}_{n,\chi,\zeta}\left(-q\right)\frac{t^{n}}{n!} &= \left[2\right]_{q}\sum_{l=0}^{d-1}\left(-1\right)^{l}q^{d-l+1}\zeta^{l}\chi\left(l\right)\frac{e^{-l(1+q)t}}{\zeta^{d}e^{-d(1+q)t}+q^{d}}\\ &= \left[2\right]_{q}\sum_{l=0}^{d-1}\left(-1\right)^{l}q^{-l+1}\zeta^{l}\chi\left(l\right)e^{-l(1+q)t}\sum_{m=0}^{\infty}\left(-1\right)^{m}\zeta^{md}q^{-md}e^{-md(1+q)t}\\ &= q\left[2\right]_{q}\sum_{m=0}^{\infty}\sum_{l=0}^{d-1}\left(-1\right)^{l+md}\chi\left(l+md\right)q^{-(l+md)}\zeta^{l+md}e^{-(l+md)(1+q)t}\\ &= q\left[2\right]_{q}\sum_{m=0}^{\infty}\left(-1\right)^{m}\zeta^{m}\chi\left(m\right)q^{-m}e^{-m(1+q)t}.\end{aligned}$$

Thus, we get the following theorem.

Theorem 2.3. *For $\zeta\in T_p$, then we have*

$$\mathcal{G}_{q,\zeta}\left(t\mid\chi\right)=\sum_{n=0}^{\infty}\mathcal{A}_{n,\chi,\zeta}\left(-q\right)\frac{t^{n}}{n!}=\left[2\right]_{q}\sum_{m=0}^{\infty}\frac{\left(-1\right)^{m}\zeta^{m}\chi\left(m\right)e^{-m(1+q)t}}{q^{m-1}}. \tag{13}$$

By using Taylor expansion of $e^{-m(1+q)t}$ in (13), we discover the following.

Theorem 2.4. *For $\zeta\in T_p$, then we have*

$$\frac{\left(-1\right)^{n}}{q\left(1+q\right)^{n+1}}\mathcal{A}_{n,\chi,\zeta}\left(-q\right)=\sum_{m=1}^{\infty}\frac{\left(-1\right)^{m}\zeta^{m}\chi\left(m\right)m^{n}}{q^{m}}. \tag{14}$$

From expressions of (12) and (14), we easily effect the following corollary:

Corollary 2.5. *For $\zeta\in T_p$ and $n\in\mathbb{N}$, then we procure the following*

$$\lim_{n\to\infty}\sum_{x=0}^{p^{n}-1}\frac{\left(-1\right)^{x}\zeta^{x}\chi\left(x\right)x^{n}}{q^{x}}=2q^{2}\sum_{m=1}^{\infty}\frac{\left(-1\right)^{m}\zeta^{m}\chi\left(m\right)m^{n}}{q^{m}}.$$

Now, we want to show multiplication theorem for Dirichlet's type of twisted Eulerian polynomials by using p-adic fermionic q-integral on $\mathbb{Z}_p$, as follows:

For each $\zeta \in \mathcal{T}_p$ and $n \in \mathbb{N}^*$, so we have

$$\begin{aligned}
&I_{-q^{-1}}\left(\zeta^x \chi(x) x^n\right) \\
&= \int_{\mathbb{Z}_p} \zeta^x \chi(x) x^n d\mu_{-q^{-1}}(x) \\
&= \lim_{m\to\infty} \frac{1}{[dp^m]_{-q^{-1}}} \sum_{x=0}^{dp^m-1} (-1)^x \zeta^x \chi(x) x^n q^{-x} \\
&= \frac{d^n}{[d]_{-q^{-1}}} \sum_{a=0}^{d-1} (-1)^a \zeta^a \chi(a) q^{-a} \left(\lim_{m\to\infty} \frac{1}{[p^m]_{-q^{-d}}} \sum_{x=0}^{p^m-1} (-1)^x \zeta^{dx} \left(\frac{a}{d}+x\right)^n q^{-dx} \right) \\
&= \frac{d^n}{[d]_{-q^{-1}}} \sum_{a=0}^{d-1} (-1)^a \chi(a) \zeta^a q^{-a} \int_{\mathbb{Z}_p} \left(\frac{a}{d}+x\right)^n \zeta^{dx} d\mu_{-q^{-d}}(x).
\end{aligned}$$

Then, we can express Dirichlet's type of twisted Eulerian polynomials in terms of p-adic fermionic q-integral on $\mathbb{Z}_p$, as follows:

Theorem 2.6. *The following*
(15)

$$\frac{(-1)^n}{(1+q)^n} \mathcal{A}_{n,\chi,\zeta}(-q) = \frac{d^n}{[d]_{-q^{-1}}} \sum_{a=0}^{d-1} (-1)^a \chi(a) \zeta^a q^{-a} \int_{\mathbb{Z}_p} \left(\frac{a}{d}+x\right)^n \zeta^{dx} d\mu_{-q^{-d}}(x)$$

is true.

Equation (15) seems to be interesting for evaluating at $q = 1$, so we see that

$$(16) \quad \frac{(-1)^n}{2^n} \mathcal{A}_{n,\chi,\zeta}(-1) = d^n \sum_{a=0}^{d-1} (-1)^a \chi(a) \zeta^a \int_{\mathbb{Z}_p} \left(\frac{a}{d}+x\right)^n \zeta^{dx} d\mu_{-1}(x).$$

Next, with the help of Rim and Kim's paper [12], we can write the following:

$$(17) \quad E_{n,\zeta}(x) = \int_{\mathbb{Z}_p} \zeta^y (x+y)^n d\mu_{-1}(y)$$

where $\zeta \in \mathcal{T}_p$ and taking $x = 0$ in the above equation, we have $E_{n,\zeta}(0) := E_{n,\zeta}$, which is defined via the following generating function:

$$(18) \quad \sum_{n=0}^{\infty} E_{n,\zeta} \frac{t^n}{n!} = \frac{2\sum_{l=0}^{d-1} (-1)^l \zeta^l e^{lt}}{\zeta^d e^{dt}+1}, \quad |t| < \frac{\pi}{d}.$$

By expressions of (16), (17) and (18), we state the following corollary.

Corollary 2.7. *For any $\zeta \in \mathcal{T}_p$ and $n \in \mathbb{N}^*$, then we obtain*

$$\mathcal{A}_{n,\chi,\zeta}(-1) = (-2d)^n \sum_{a=0}^{d-1} (-1)^a \chi(a) \zeta^a E_{n,\zeta^d}\left(\frac{a}{d}\right).$$

3. On the twisted Eulerian-$\mathcal{L}$-function in $\mathbb{C}$

The objective of this part is to describe twisted Eulerian-$\mathcal{L}$-function by applying Mellin transformation to the generating function of Dirichlet's type of twisted Eulerian polynomials, which seem to be interpolation function of Dirichlet's type of twisted Eulerian polynomials at negative integers by using Cauchy-Residue theorem. We want to determine that these functions are very useful in Number Theory, Complex Analysis and Mathematical Physics, p-adic analysis and other areas cf. [1], [4], [5], [7], [11], [14], [15], [17], [18], [20], [24], [25] and [26]. Thus, by (13), for $s \in \mathbb{C}$, we consider the following integral:

$$\mathcal{L}_{E,\zeta}(s,\chi) = \frac{\int_0^\infty t^{s-1}\mathcal{G}_{q,\zeta}(t \mid \chi)\,dt}{\int_0^\infty t^{s-1}e^{-t}dt}. \tag{19}$$

By (19), we can derive the following:

$$\begin{aligned}\mathcal{L}_{E,\zeta}(s,\chi) &= q[2]_q \sum_{m=0}^{\infty} (-1)^m \chi(m) \zeta^m q^{-m} \left\{ \frac{\int_0^\infty t^{s-1}e^{-m(1+q)t}dt}{\int_0^\infty t^{s-1}e^{-t}dt} \right\} \\ &= \frac{q}{(1+q)^{s-1}} \sum_{m=1}^{\infty} \frac{(-1)^m \chi(m) \zeta^m}{q^m m^s}.\end{aligned}$$

On the other hand, we easily see that

$$= \sum_{n=0}^{\infty} \frac{\mathcal{A}_{n,\chi,\zeta}(-q)}{n!} \left(\frac{\int_0^\infty t^{s-n+1}dt}{\int_0^\infty t^{s-1}e^{-t}dt} \right). \tag{20}$$

So, we give definition of twisted Eulerian $\mathcal{L}$-function as follows:

Definition 2. *Let $\zeta \in \mathcal{T}_p$ and $s \in \mathbb{C}$, then we define the following*

$$\mathcal{L}_{E,\zeta}(s,\chi) = \frac{q}{(1+q)^{s-1}} \sum_{m=1}^{\infty} \frac{(-1)^m \chi(m) \zeta^m}{q^m m^s}.$$

By using (19) and (20), then, we derive the following which seems to be interesting and useful for improving of Complex Analysis and Theory of Analytic Numbers.

Theorem 3.1. *The following equality holds true:*

$$\mathcal{L}_{E,\zeta}(-n,\chi) = (-1)^n \mathcal{A}_{n,\chi,\zeta}(-q).$$

References

[1] E. Cetin, M. Acikgoz, I. N. Cangul and S. Araci, A note on the (h,q)-Zeta type function with weight α, http://arxiv.org/pdf/1206.5299.pdf.

[2] D. S. Kim, T. Kim, W. J. Kim and D. V. Dolgy, A note on Eulerian polynomials, *Abstract and Applied Analysis*, Volume 2012 (2012), Article ID 269640, 10 pages.

[3] T. Kim, On a q-analogue of the p-adic log gamma functions and related integrals, *J. Number Theory* 76 (1999), 320-329.

[4] T. Kim, q-Volkenborn integration, Russ. J. Math Phys., 19 (2002), 288-299.

[5] T. Kim, Non-archimedean q-integrals associated with multiple Changhee q-Bernoulli Polynomials, *Russ. J. Math Phys.*, 10 (2003), 91-98.

[6] T. Kim, p-adic q-integrals associated with the Changhee-Barnes' q-Bernoulli Polynomials, *Integral Transform. Spec. Funct.*, 15 (2004), 415-420.

[7] T. Kim, Analytic continuation of multiple q-zeta functions and their values at negative integers, *Russ. J. Math Phys.*, 11 (2004), 71-76.

[8] T. Kim, Some identities on the q-Euler polynomials of higher order and q-stirling numbers by the fermionic p-adic integral on $\mathbb{Z}_p$, *Russian J. Math. Phys.* 16 (2009), 484–491.

[9] T. Kim, On the q-extension of Euler and Genocchi numbers, *J. Math. Anal. Appl.* 326 (2007) 1458–1465.

[10] T. Kim, On the analogs of Euler numbers and polynomials associated with p-adic q-integral on $\mathbb{Z}_p$ at $q = 1$, *J. Math. Anal. Appl.* 331 (2007) 779–792.

[11] T. Kim, S. H. Lee, H. H. Han and C. S. Ryoo, On the values of the weighted q-Zeta and L-functions, *Discrete Dynamics in Nature and Society*, Volume 2011, Article ID 476381, 7 pp.

[12] S.-H. Rim and T. Kim, A note on q-Euler numbers associated with the basic q-zeta function, *Applied Mathematics Letters*, vol. 20, no. 4, pp. 366–369, 2007.

[13] D. Foata, Eulerian polynomials: from Euler's time to the present, *The legacy of Alladi Ramakrishnan in the mathematical sciences*, 253–273, Springer, New York, 2010.

[14] L. C. Jang, The q-analogue of twisted Lerch type Euler Zeta functions, *Bull. Korean Math. Soc.* 47 (2010), No. 6, pp. 1181-1188.

[15] L. C. Jang, V. Kurt, Y. Simsek, and S. H. Rim, q-analogue of the p-adic twisted l-function, *Journal of Concrete and Applicable Mathematics*, vol. 6, no. 2, pp. 169–176, 2008.

[16] H. Ozden, I. N. Cangul, Y. Simsek, Multivariate interpolation functions of higher order q-Euler numbers and their applications, *Abstract and Applied Analysis* 2008 (2008), Article ID 390857, 16 pages.

[17] I. N. Cangul, H. Ozden, Y. Simsek, A new approach to q-Genocchi numbers and their interpolation functions, *Nonlinear Analysis* 71 (2009), pp. 793-799.

[18] I. N. Cangul, H. Ozden, and Y. Simsek, Generating functions of the (h, q) extension of twisted Euler polynomials and numbers, *Acta Mathematica Hungarica*, vol. 120, no. 3, pp. 281–299, 2008.

[19] M. Acikgoz and Y. Simsek, On Multiple Interpolation functions oof the Nörlund-Type q-Euler polynomials, *Abstract and Applied Analysis*, Volume 2009, Article ID 382574, 14 pages.

[20] S. Araci, M. Acikgoz, K. H. Park and H. Jolany, On the unification of two families of multiple twisted type polynomials by using p-adic q-integral on $\mathbb{Z}_p$ at $q = -1$, *Bulletin of the Malaysian Mathematical Sciences and Society* (accepted for publication).

[21] S. Araci, M. Acikgoz and H. Jolany, On p-adic interpolation function associated with modified Dirichlet's type of twisted q-Euler numbers and polynomials with weight α, http://arxiv.org/pdf/1201.5490v1.pdf.

[22] S. Araci, M. Acikgoz and D. Gao, On the Dirichlet's type of Eulerian polynomials, http://arxiv.org/pdf/1207.1834.pdf.

[23] S. Araci, D. Erdal and J. J. Seo, A study on the fermionic p-adic q-integral representation on $\mathbb{Z}_p$ associated with weighted q-Bernstein and q-Genocchi polynomials, *Abstract and Applied Analysis*, Volume 2011, Article ID 649248, 10 pages.

[24] Y. Simsek, q-analogue of twisted l-series and q-twisted Euler numbers, *Journal of Number Theory*, vol. 110, no. 2, pp. 267–278, 2005.

[25] Y. Simsek, q-Dedekind type sums related to q-zeta function and basic L-series, *Journal of Mathematical Analysis and Applications*, vol. 318, no. 1, pp. 333–351, 2006.

[26] Y. Simsek, Twisted (h, q)-Bernoulli numbers and polynomials related to twisted (h, q)-zeta function and L-function, *Journal of Mathematical Analysis and Applications*, vol. 324, no. 2, pp. 790–804, 2006.

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics, 27310 Gaziantep, TURKEY
E-mail address: `mtsrkn@hotmail.com`

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics, 27310 Gaziantep, TURKEY
E-mail address: `acikgoz@gantep.edu.tr`